arXiv:math/0406323v1 [math.CO] 16 Jun 2004

Some Formulae for Bivariate Fibonacci and Lucas Polynomials

Mario Catalani
Department of Economics, University of Torino
Via Po 53, 10124 Torino, Italy
mario.catalani@unito.it

Abstract

We derive a collection of identities for bivariate Fibonacci and Lucas polynomials using essentially a matrix approach as well as properties of such polynomials when the variables x and y are replaced by polynomials. A wealth of combinatorial identities can be obtained for selected values of the variables.

1 Introduction

We define bivariate Fibonacci polynomials as

$$F_n(x,\,y) = xF_{n-1}(x,\,y) + yF_{n-2}(x,\,y), \qquad F_0(x,\,y) = 0,\ F_1(x,\,y) = 1,$$

and bivariate Lucas polynomials as

$$L_n(x,\,y) = xL_{n-1}(x,\,y) + yL_{n-2}(x,\,y), \qquad L_0(x,\,y) = 2,\ L_1(x,\,y) = x.$$

We assume $x \neq 0,\ y \neq 0,\ x^2 + 4y \neq 0$.
The roots of the characteristic equation are

$$\alpha \equiv \alpha(x,\,y) = \frac{x + \sqrt{x^2+4y}}{2}, \qquad \beta \equiv \beta(x,\,y) = \frac{x - \sqrt{x^2+4y}}{2}.$$

We have $\alpha + \beta = x$, $\alpha\beta = -y$ and $\alpha - \beta = \sqrt{x^2+4y}$. The Binet's forms are

$$F_n(x,\,y) = \frac{\alpha^n - \beta^n}{\alpha - \beta}, \qquad L_n(x,\,y) = \alpha^n + \beta^n.$$

Many basic facts concerning these kinds of polynomials can be found in [1].

2 Some Matrix Techniques

Let

$$\mathbf{B} = \left[\begin{array}{cc} x^2 + 2y & x \\ xy & 2y \end{array} \right].$$

Then

$$\mathbf{B} = 2y\mathbf{I} + x\mathbf{A},$$

where

$$\mathbf{A} = \left[\begin{array}{cc} x & 1 \\ y & 0 \end{array} \right].$$

It follows

$$\mathbf{B}^n = \sum_{k=0}^{n} \binom{n}{k} (2y)^{n-k} x^k \mathbf{A}^k. \tag{1}$$

It is known that (e.g. [1])

$$\mathbf{A}^n = \left[\begin{array}{cc} F_{n+1}(x,\, y) & F_n(x,\, y) \\ yF_n(x,\, y) & yF_{n-1}(x,\, y) \end{array} \right].$$

Now using the method exposed in [2] to $\mathbf{B}$ we find, after some manipulations, that the element $(1,\, 2)$ of $\mathbf{B}^n$ is given by

$$x \sum_{k=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n-1-k}{k} (x^2 + 4y)^{n-1-k} (-y)^k.$$

The corresponding element of the RHS of Equation 1 is

$$\sum_{k=0}^{n} \binom{n}{k} (2y)^{n-k} x^k F_k(x,\, y).$$

Then, using the Binet's form,

$$x \sum_{k=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n-1-k}{k} (x^2+4y)^{n-1-k} (-y)^k = \frac{1}{\alpha - \beta} \sum_{k=0}^{n} \binom{n}{k} (2y)^{n-k} x^k (\alpha^k - \beta^k).$$

Now

$$\begin{aligned} \sum_{k=0}^{n} \binom{n}{k} (2y)^{n-k} x^k \alpha^k &= (2y + x\alpha)^n \\ &= \alpha^n (\alpha - \beta)^n. \end{aligned}$$

Analogously for the β part. After replacing n by $2n$ we get the identity

$$x\sum_{k=0}^{n-1}\binom{2n-1-k}{k}(x^2+4y)^{n-k-1}(-y)^k = F_{2n}(x,\,y). \tag{2}$$

Now

$$\mathbf{BA} = \left[\begin{array}{cc} x^3+3xy & x^2+2y \\ x^2y+2y^2 & xy \end{array}\right],$$

and

$$\begin{gathered}\operatorname{tr}(\mathbf{BA}) = x^3+4xy = x(x^2+4y), \\ |\mathbf{BA}| = |\mathbf{B}|\,|\mathbf{A}| = -y(x^2y+4y^2) = -y^2(x^2+4y).\end{gathered}$$

Applying the previous method this time to $\mathbf{BA}$ we get that the element $(1,\,2)$ of $(\mathbf{BA})^n$ is given by

$$\begin{aligned}(x^2+2y)\sum_{k=0}^{\lfloor\frac{n-1}{2}\rfloor}\binom{n-1-k}{k}x^{n-1-2k}(x^2+4y)^{n-1-2k}y^{2k}(x^2+4y)^k = \\ = (x^2+2y)\sum_{k=0}^{\lfloor\frac{n-1}{2}\rfloor}\binom{n-1-k}{k}x^{n-1-2k}(x^2+4y)^{n-1-k}y^{2k}.\end{aligned}$$

On the other hand (see [1])

$$\mathbf{BA} = (2y\mathbf{I}+x\mathbf{A})\mathbf{A} = 2y\mathbf{A}+x\mathbf{A}^2,$$

so that

$$\begin{aligned}(\mathbf{BA})^n &= \sum_{k=0}^{n}\binom{n}{k}(2y)^{n-k}\mathbf{A}^{n-k}x^k\mathbf{A}^{2k} \\ &= \sum_{k=0}^{n}\binom{n}{k}(2y)^{n-k}x^k\mathbf{A}^{n+k}.\end{aligned}$$

Since the element $(1,\,2)$ of $\mathbf{A}^n$ is $F_n(x,\,y)$, we have that the element $(1,\,2)$ of $(\mathbf{BA})^n$ is

$$\sum_{k=0}^{n}\binom{n}{k}(2y)^{n-k}x^kF_{n+k}(x,\,y) = \frac{1}{\alpha-\beta}\sum_{k=0}^{n}\binom{n}{k}(2y)^{n-k}x^k(\alpha^{n+k}-\beta^{n+k}).$$

Now

$$\begin{aligned}
\frac{1}{\alpha-\beta}\sum_{k=0}^{n}\binom{n}{k}(2y)^{n-k}x^k\alpha^{n+k} &= \frac{\alpha^n}{\alpha-\beta}\sum_{k=0}^{n}\binom{n}{k}(2y)^{n-k}(x\alpha)^k \\
&= \frac{\alpha^n}{\alpha-\beta}(2y+x\alpha)^n \\
&= \frac{\alpha^n}{\alpha-\beta}\alpha^n(\alpha-\beta)^n \\
&= \frac{\alpha^{2n}(\alpha-\beta)^n}{\alpha-\beta}.
\end{aligned}$$

In the same way

$$\begin{aligned}
\frac{1}{\alpha-\beta}\sum_{k=0}^{n}\binom{n}{k}(2y)^{n-k}x^k\alpha^{n+k} &= \frac{\beta^n}{\alpha-\beta}\sum_{k=0}^{n}\binom{n}{k}(2y)^{n-k}(x\beta)^k \\
&= \frac{\beta^n}{\alpha-\beta}(2y+x\beta)^n \\
&= \frac{\beta^n}{\alpha-\beta}\beta^n(-1)^n(\alpha-\beta)^n \\
&= \frac{\beta^{2n}(-1)^n(\alpha-\beta)^n}{\alpha-\beta}.
\end{aligned}$$

Subtracting we get

$$(\alpha-\beta)^n\left(\frac{\alpha^{2n}}{\alpha-\beta}-\frac{\beta^{2n}(-1)^n}{\alpha-\beta}\right).$$

Replacing n with $2n$ and equating the two expressions for the element (1, 2) we get

$$\begin{aligned}
&(x^2+2y)\sum_{k=0}^{n-1}\binom{2n-1-k}{k}x^{2n-1-2k}(x^2+4y)^{2n-1-k}y^{2k} \\
&\quad=(x^2+4y)^nF_{4n}(x,\,y),
\end{aligned}$$

that is

$$(x^2+2y)\sum_{k=0}^{n-1}\binom{2n-1-k}{k}x^{2n-1-2k}(x^2+4y)^{n-1-k}y^{2k}=F_{4n}(x,\,y). \tag{3}$$

Using Equation 2 and replacing here n by $2n$ we obtain the identity

$$\begin{aligned}&(x^2+2y)\sum_{k=0}^{n-1}\binom{2n-1-k}{k}x^{2n-1-2k}(x^2+4y)^{n-1-k}y^{2k}\\&=x\sum_{k=0}^{2n-1}\binom{4n-1-k}{k}(x^2+4y)^{2n-1-k}(-y)^k.\end{aligned}$$

3 Polynomials as Arguments

First of all, from

$$F_n(x,\,y)=\frac{\alpha^n-\beta^n}{\alpha-\beta},\qquad L_n(x,\,y)=\alpha^n+\beta^n,$$

solving for α and β we get

$$\alpha^n=\frac{L_n(x,\,y)+(\alpha-\beta)F_n(x,\,y)}{2},\tag{4}$$

$$\beta^n=\frac{L_n(x,\,y)-(\alpha-\beta)F_n(x,\,y)}{2}.\tag{5}$$

Secondly

$$\begin{aligned}L_n^2(x,\,y)&=\alpha^{2n}+\beta^{2n}+2(-1)^ny^n\\&=\alpha^{2n}+\beta^{2n}-2(-1)^{n+1}y^n,\end{aligned}$$

$$\begin{aligned}(\alpha-\beta)^2F_n^2(x,\,y)&=\alpha^{2n}+\beta^{2n}-2(-1)^ny^n\\&=\alpha^{2n}+\beta^{2n}+2(-1)^{n+1}y^2.\end{aligned}$$

It follows

$$L_n^2(x,\,y)+(-1)^{n+1}4y^n=(x^2+4y)F_n^2(x,\,y).\tag{6}$$

Then

$$\begin{aligned}\alpha\left(L_n(x,\,y),\,(-1)^{n+1}y^n\right)&=\frac{L_n(x,\,y)+\sqrt{L_n^2(x,\,y)+4(-1)^{n+1}y^n}}{2}\\&=\frac{L_n(x,\,y)+\sqrt{x^2+4y}F_n(x,\,y)}{2}\\&=\alpha^n(x,\,y).\end{aligned}$$

$$\beta\left(L_n(x,\,y),\,(-1)^{n+1}y^n\right) = \beta^n(x,\,y).$$

Then

$$\begin{aligned}
&F_n\left(L_k(x,\,y),\,(-1)^{k+1}y^k\right)\\
&\quad= \frac{\alpha^n\left(L_k(x,\,y),\,(-1)^{k+1}y^k\right) - \beta^n\left(L_k(x,\,y),\,(-10^{k+1}y^k\right)}{\alpha\left(L_k(x,\,y),\,(-1)^{k+1}y^k\right) - \beta\left(L_k(x,\,y),\,(-1)^{k+1}y^k\right)}\\
&\quad= \frac{\alpha^{nk}(x,\,y) - \beta^{nk}(x,\,y)}{\alpha^k(x,\,y) - \beta^k(x,\,y)}\\
&\quad= \frac{F_{nk}(x,\,y)}{F_k(x,\,y)}. \qquad (7)
\end{aligned}$$

As a consequence, we can see that $F_k(x,\,y)$ divides $F_{kn}(x,\,y)$, $\forall\,k,\,n$. The analogous of Equation 7 for Lucas polynomials is

$$L_n\left(L_k(x,\,y),\,(-1)^{k+1}y^k\right) = L_{nk}(x,\,y). \qquad (8)$$

Some examples.

$$F_{2n}(x,\,y) = F_2(x,\,y)F_n(L_2(x,\,y),\,-y^2) = xF_n(x^2+2y,\,-y^2),$$

$$F_{3n}(x,\,y) = F_3(x,\,y)F_n(L_3(x,\,y),\,y^3) = (x^2+y)F_n(x^3+3xy,\,y^3),$$

$$F_{4n}(x,\,y) = F_4(x,\,y)F_n(L_4(x,\,y),\,-y^4) = x(x^2+2y)F_n(x^4+4x^2y+2y^2,\,-y^4).$$

Equation 6 and Equation 8 can be used to derive many other identities. For instance the *Simpson* formula (see [1])

$$L_n(x,\,y)L_{n+2}(x,\,y) - L^2_{n+1}(x,\,y) = (-1)^n y^n(x^2+4y) \qquad (9)$$

setting $x = L_k(x,\,y)$, $y = (-1)^{k+1}y^k$ becomes

$$\begin{aligned}
&L_n\left(L_k(x,\,y),\,(-1)^{k+1}y^k\right)L_{n+2}\left(L_k(x,\,y),\,(-1)^{k+1}y^k\right) +\\
&\qquad -L^2_{n+1}\left(L_k(x,\,y),\,(-1)^{k+1}y^k\right)\\
&\quad= (-1)^n((-1)^{k+1}y^k)^n(L^2_k(x,\,y) + 4(-1)^{k+1}y^k)\\
&\quad= (-y)^{nk}(x^2+4y)F^2_k(x,\,y),
\end{aligned}$$

that is

$$L_{kn}(x,\, y)L_{k(n+2)}(x,\, y) - L^2_{k(n+1)}(x,\, y) = (-y)^{nk}(x^2 + 4y)F^2_k(x,\, y).$$

Equation 2 with $x = L_k(x,\, y)$, $y = (-1)^{k+1}y^k$ becomes

$$\begin{aligned} L_k(x,\, y)\sum_{r=0}^{n-1}\binom{2n-1-r}{r}(x^2 + 4y)^{n-1-r}F_k^{2(n-1-r)}(x,\, y)(-y)^{rk} \\ = F_{2n}(L_k(x,\, y),\, (-1)^{k+1}y^k). \end{aligned}$$

Now using the identity

$$L^2_n(x,\, y) + 2(-1)^{n+1}y^n = L_{2n}(x,\, y),$$

we have

$$\begin{aligned} &F_{2n}\left(L_k(x,\, y),\, (-1)^{k+1}y^k\right) \\ &\quad = L_k(x,\, y)F_n\left(L^2_k(x,\, y) + 2(-1)^{k+1}y^k,\, -((-1)^{k+1}y^k)^2\right) \\ &\quad = L_k(x,\, y)F_n\left(L_{2k}(x,\, y),\, -y^{2k}\right). \end{aligned}$$

Finally

$$\sum_{r=0}^{n-1}\binom{2n-1-r}{r}(x^2+4y)^{n-1-r}F_k^{2(n-1-r)}(x,\, y)(-y)^{rk} = F_n\left(L_{2k}(x,\, y),\, -y^{2k}\right).$$

Then, since $-y^{2k} = (-1)^{2k+1}y^{2k}$, Equation 7 allows to write

$$\sum_{r=0}^{n-1}\binom{2n-1-r}{r}(x^2 + 4y)^{n-1-r}F_k^{2(n-1-r)}(x,\, y)(-y)^{rk} = \frac{F_{2kn}(x,\, y)}{F_{2k}(x,\, y)}.$$

Equation 7 and Equation 8 allow to derive new identities. For example, the identity

$$yF_{n-1}(x,\, y) + F_{n+1}(x,\, y) = L_n(x,\, y) \tag{10}$$

upon substitution of x with $L_k(x,\, y)$ and y with $(-1)^{k+1}y^k$ becomes

$$(-1)^{k+1}y^k\frac{F_{k(n-1)}(x,\, y)}{F_k(x,\, y)} + \frac{F_{k(n+1)}(x,\, y)}{F_k(x,\, y)} = L_{nk}(x,\, y),$$

that is

$$(-1)^{k+1}y^kF_{k(n-1)}(x,\, y) + F_{k(n+1)}(x,\, y) = F_k(x,\, y)L_{nk}(x,\, y).$$

Or the identity

$$L_{n+2}^2(x,\,y)+yL_{n+1}^2(x,\,y)=(x^2+2y)L_{2n+2}(x,\,y)+xyL_{2n+1}(x,\,y)$$

that becomes

$$\begin{aligned}&L_{k(n+2)}^2(x,\,y)+(-1)^{k+1}y^kL_{k(n+1)}^2(x,\,y)\\&\qquad=L_{2k}(x,\,y)L_{k(2n+2)}(x,\,y)+(-1)^{k+1}y^kL_k(x,\,y)L_{k(2n+1)}(x,\,y).\end{aligned}$$

Equation 6 can be rewritten as

$$L_n^2(x,\,y)=(x^2+4y)F_n^2(x,\,y)+(-1)^n4y^n.$$

Then

$$\alpha\left(\sqrt{x^2+4y}F_n(x,\,y),\,(-1)^ny^n\right)=\alpha^n(x,\,y),$$

$$\beta\left(\sqrt{x^2+4y}F_n(x,\,y),\,(-1)^ny^n\right)=-\beta^n(x,\,y).$$

It follows

$$F_{2n+1}\left(\sqrt{x^2+4y}F_k(x,\,y),\,(-1)^ky^k\right)=\frac{L_{k(2n+1)}(x,\,y)}{L_k(x,\,y)},\tag{11}$$

$$F_{2n}\left(\sqrt{x^2+4y}F_k(x,\,y),\,(-1)^ky^k\right)=(\alpha-\beta)\frac{F_{2kn}(x,\,y)}{L_k(x,\,y)}.\tag{12}$$

For the Lucas polynomials we have

$$L_{2n+1}\left(\sqrt{x^2+4y}F_k(x,\,y),\,(-1)^ky^k\right)=(\alpha-\beta)F_{k(2n+1)}(x,\,y),\tag{13}$$

$$L_{2n}\left(\sqrt{x^2+4y}F_k(x,\,y),\,(-1)^ky^k\right)=L_{2kn}(x,\,y).\tag{14}$$

Now identity 10 replacing x with $\sqrt{x^2+4y}F_k(x,\,y)$ and y with $(-1)^ky^k$ becomes, using Equations 11, 12, 13, 14,

$$(-1)^ky^kL_{k(2n-1)}(x,\,y)+L_{k(2n+1)}(x,\,y)=L_{2kn}(x,\,y)L_k(x,\,y),$$

and

$$(-1)^ky^kF_{2kn}(x,\,y)+F_{k(2n+2)}(x,\,y)=F_{k(2n+1)}(x,\,y)L_k(x,\,y).$$

In an analogous way the Simpson formula (Equation 9) becomes

$$L_{2kn}(x,\,y)L_{k(2n+2)}(x,\,y)-(\alpha-\beta)^2F_{k(2n+1)}^2(x,\,y)=y^{2nk}L_k^2(x,\,y),$$

and

$$(\alpha-\beta)^2F_{k(2n-1)}(x,\,y)F_{k(2n+1)}(x,\,y)-L_{2kn}^2(x,\,y)=-y^{k(2n-1)}L_k^2(x,\,y).$$

References

[1] M. Catalani (2004), ”Generalized Bivariate Fibonacci Polynomials.” Version 2. http://front.math.ucdavis.edu/math.CO/0211366

[2] J. Mc Laughlin (2004), ”Combinatorial Identities Deriving from the n-th Power of a 2×2 Matrix”, published electronically at http://emp.trincoll.edu/~jmclaugh/2x2atrixpowerglaintegers.pdf